

## Estimate of permafrost organic carbon balance in Alaskan boreal and tundra ecosystems using natural level radiocarbon

Miyuki Kondo<sup>1</sup>, Masao Uchida<sup>1</sup>, Motoo Utsumi<sup>2</sup>, Go Iwahana<sup>3</sup>, Hiroki Iwata<sup>4</sup>, Yoshinobu Harazono<sup>3</sup>, Taro Nakai<sup>3,5</sup>, Kiyoshi Tanabe<sup>1</sup>, and Yasuyuki Shibata<sup>1</sup>

<sup>1</sup> *National Institute for Environmental Studies*

<sup>2</sup> *University of Tsukuba*

<sup>3</sup> *International Arctic Research Center (IARC), University of Alaska Fairbanks*

<sup>4</sup> *Kyoto University*

<sup>5</sup> *Researcher at Hydrospheric Atmospheric Research Center, Nagoya University*

The high-latitude regions, where a serious warming is expected, currently store large amounts of soil organic carbon (SOC) in active-layer soils and permafrost, accounting for nearly half of the global belowground organic carbon pool. Despite the importance of these regions in the present carbon cycle, the soil C fluxes and budget are still only poorly known. Here, we use radiocarbon as the tool for quantifying the C balance of the inputs and decomposition in tundra and boreal soil. We evaluated the C inputs (I) and decomposition rates (k, inverse of turnover time) and net C accumulation (CA), using <sup>14</sup>C approaches.

Tundra and boreal soils show different patterns of depth distribution and C storage. Cumulative organic carbon stocks in boreal forest are 5.3 and 19.2 kgCm<sup>-2</sup>, in surface organic layer (0-25 cm), and deep organic and mineral layers (25-70 cm), respectively. Large annual C input (0.249 kgCm<sup>-2</sup> yr<sup>-1</sup>) and relatively slow decomposition (27 years) lead to rapid CA (0.052 kgCm<sup>-2</sup> yr<sup>-1</sup>) in surface organic layer in boreal forest. Deep organic and mineral layers including near-surface permafrost show slower rate of input (0.031 kgCm<sup>-2</sup> yr<sup>-1</sup>) and turnover (617 years) and CA about 20 times slower (0.003 kgCm<sup>-2</sup> yr<sup>-1</sup>) than surface organic layer. Decomposition organic matter (Rh), which in accord with C losses from both surface and subsurface layers, was 0.225 kgCm<sup>-2</sup> yr<sup>-1</sup>. This value agreed well with Rh (0.227 kgCm<sup>-2</sup> yr<sup>-1</sup>) simulated by process-based models that simulate the biogeochemical and hydrologic cycle, where Rh averaged 45% of ecosystem respiration and 59% of soil respiration<sup>2</sup>. In contrast, large amount of SOC (36.4 kg m<sup>-2</sup>) have accumulated over millennia (turnover time: 4540 yrs) below the thin organic layer in tundra. The CA of mineral layer and permafrost is close to zero (0.003 kgCm<sup>-2</sup> yr<sup>-1</sup>), and Rh is 0.008 kgCm<sup>-2</sup> yr<sup>-1</sup>. Our radiocarbon data show that the most SOC in tundra soil was mode of stabilizing OC by permafrost and steady-state SOC stocks under current C balance.

### References

1. Ueyama M., Y. Harazono, Y. Kim and N.Tanaka, Response of the carbon cycle in sub-arctic black spruce forests to climate change: Reduction of a carbon sink related to the sensitivity of heterotrophic respiration, Agric. For. Meteorol., 149, 582–602, 2009.